Chemistry 30 Unit 4: Solutions Assignment 4 - The Solubility Product Constant, K_{sp}

1. Write the balanced equation **and** the solubility product constant expression, K_{sp}, for the each of the following dissociation reactions. All compounds are solids. One has been given as an example.

An important reminder – the seven diatomic molecules $(H_2, N_2, O_2, F_2, Cl_2, Br_2, I_2)$ **ARE NOT** diatomic as ions!

Reminders – ion charges MUST BE included.

- solids (and liquids) are NOT included in the equilibrium expression
- don't forget to include exponents when needed
- polyatomic ions (e.g. CO₃) do not break apart

Compound	Equation	K _{sp}
(NH ₄) ₂ S	$(NH_4)_2O(s) = 2 NH_4^+(aq) + S^{2-}(aq)$	K _{sp} = [NH ₄ ⁺] ² [S ²⁻]
CaS		
K ₂ SO ₄		
Mg(OH) ₂		

2. Consider the K_{sp} values for the following substances, all measured at 25°C

PbCrO ₄	2.0×10^{-16}
PbSO ₄	1.3 × 10⁻ ⁸

PbCO₃ 7.4×10^{-14}

Which substance is MOST soluble (dissolves the best)

Which substances is LEAST soluble (dissolves the worst)

3. Solutions are prepared by dissolving a sulfate salt (an ionic compound containing the sulfate ion) in water. Which one of the compounds listed would we use in order to get the highest concentration of sulfate ion, SO₄²⁻?

(a salt is an ionic compound that contains a positive ion other than hydrogen and a negative ion other than hydroxide).

- A. $CaSO_4$ K_{sp} = 2.6×10^{-4}
- B. SrSO₄ $K_{sp} = 7.6 \times 10^{-7}$
- C. BaSO₄ $K_{sp} = 1.5 \times 10^{-9}$
- D. RaSO₄ $K_{sp} = 4.0 \times 10^{-11}$
- 4. Calculate K_{sp} for a saturated nickel(II) sulfide, NiS, solution with a molar concentration of 3.27 \times 10^{-11}

Begin with a balanced dissociation equation:

From information provided in the question and the balanced equation, determine the concentration of the ions Ni^{2+} and S^{2-} :

Write the K_{sp} expression for this reaction:

Substitute values into the equation and solve for the unknown, K_{sp}:

5. At 25°C, the concentration of Ce(OH)₃ in a saturated solution is 5.1×10^{-6} M. Calculate $K_{sp}.$

Begin with a balanced equation:

Determine the concentration of the ions in the solution. Be careful with [OH⁻].

Write the K_{sp} expression, then substitute values into the equation and solve for the unknown, $K_{\text{sp}}.$

6. Calculate the concentration of ions in a saturated solution of CaCO₃ in water at 25°C. K_{sp} for CaCO₃ is 4.8×10^{-9} .